

Aquatics and Marine Ecosystems

CULVERT TEST BED: FISH-PASSAGE RESEARCH FACILITY

Walter Pearson (Phone: 360-681-3661, Email: walter.pearson@pnl.gov), Associate Director, and
Christopher May (Phone: 360-681-4556, Email: christopher.may@pnl.gov), Senior Research
Engineer, Marine Sciences Laboratory, 1529 West Sequim Bay Road, Sequim, WA 98382,
Fax: 360-681-3661

Abstract

The passage of juvenile salmonids and other fish through culverts is a significant Endangered Species Act (ESA) issue throughout the Pacific Northwest and now in other areas of the nation. Much of recent research and engineering has focused on increased passage of returning adult salmon; however, juvenile-salmonid movement both up and downstream throughout the year is now recognized as substantial and is a key area in which future research promises practical returns. Because a large percentage of the culverts beneath roads in the Pacific Northwest are judged as blocking juvenile salmon from thousands of miles of habitat, determining appropriate hydraulic and fish-passage designs for retrofitted culverts before installation has both substantial cost and environmental implications.

To address these issues, the Washington State Department of Transportation (WSDOT) leads a partnership that includes the Washington Department of Fish and Wildlife (WDFW), Alaska Department of Transportation, Alaska Department of Fish and Game, Oregon Department of Transportation, California Department of Transportation, the Federal Highway Administration, and the Pacific Northwest National Laboratory (PNNL). The partnership has undertaken a phased program conducted by an interdisciplinary team of scientists and engineers from PNNL to address the hydraulic and behavioral issues associated with juvenile-salmonid fish passage through culvert systems. This program addresses the testing and assessment of full-scale physical models of culvert systems deployed in an experimental test bed. Experiments in the test bed have begun and will measure the hydraulic conditions (mean velocity, turbulence, and water depth) associated with various culvert designs under various slopes and flow regimes, and then relate these measures to repeatable, quantitative measures of fish-passage success.

The culvert test-bed program is a one-of-a-kind capability designed to provide scientifically sound information that can be used to develop better designs for retrofitted culvert installations. Compared with field studies or temporary installations, the facility promises fast results, scientific and statistically controlled evaluations, an ability to quickly discern optimum engineering principles, and elimination of expensive trial-and-error approaches of field installations.

Biographical Sketches: Dr. Walter H. Pearson is associate director of the Marine Sciences Laboratory in Sequim, Washington, which is a part of the Pacific Northwest National Laboratory, operated for the U.S. Department of Energy by Battelle Memorial Institute. His bachelor's and master's degrees are in biology from Bates College and the University of Alaska. His doctorate is in oceanography from Oregon State University. Dr. Pearson's primary area of expertise is the study of the effects of pollution and human activities on marine and estuarine environments, especially on the fisheries these environments support. He has expertise in ecotoxicology and the behavioral ecology of fish and shellfish. Working for the Marine Sciences Laboratory for over 20 years, Dr. Pearson has gained extensive experience leading large multidisciplinary, multi-organizational studies to address environmental and fisheries issues. From 1993 through 1997, Dr. Pearson was founding program director of the innovative offsite program of Huxley College of Environmental Studies in Port Angeles, Washington, as part of Western Washington University's extended education program. In 1998, Dr. Pearson joined the newly formed Environmental Research and Wildlife Development Agency (ERWDA) in Abu Dhabi in the United Arab Emirates (UAE). He served as head of the Marine Environmental Research Center (MERC) until August 2000. Dr. Pearson led MERC as its staff developed programs for sea turtles, dugong, sea grasses, fisheries, water quality, oil-spill contingency planning, coastal-sensitivity mapping, and other marine issues. He returned to the Marine Sciences Laboratory in 2000. Dr. Pearson's current research at the Marine Sciences Laboratory addresses the effects of dredging on Dungeness crab, oil-spill impacts on marine fisheries, and juvenile fish-passage through culvert systems.

Dr. Christopher May is a senior research engineer at the Marine Sciences Laboratory in Sequim, Washington, which is a part of the Pacific Northwest National Laboratory, operated for the U.S. Department of Energy by Battelle Memorial Institute. Dr. May works to extend existing ecosystem assessment and restoration capabilities in the marine and near-shore environment into freshwater ecosystems with a focus on watershed analysis, stormwater management, non-point-source pollution issues, and salmonid-habitat assessment. Dr. May has served as a researcher and adjunct faculty member at the University of Washington and Western Washington University, as a private consultant, and as a technical advisor to the U.S. Navy and Department of Defense for stormwater and watershed-management issues. Dr. May has been principal investigator on projects ranging from a study to evaluate the impacts of urbanization on aquatic ecosystems and the effectiveness of stormwater best-management practices for the Watershed Management Institute and Environmental Protection Agency to the Kitsap and Jefferson County Salmonid Refugia Projects to identify and evaluate potential salmonid habitat-conservation areas for endangered salmon. Dr. May holds a Ph.D. in environmental science and engineering from the University of Washington, an M.S. in industrial engineering and management from the University of Minnesota, and a B.S. in marine engineering from the United States Naval Academy.

ENGINEERED LOGJAMS: AN ALTERNATIVE BANK-PROTECTION METHOD FOR US 101 ALONG THE HOH RIVER, WASHINGTON

Carl Ward (Phone: 360-570-6706, Email: WardC@wsdot.wa.gov), Regional Biologist/Olympic Region Biology Program Manager, Washington State Department of Transportation, P.O. Box 47417, Olympia, WA 98504-7417, Fax: 360-570-6697

Abstract

The Washington State Department of Transportation (WSDOT) has repetitively made emergency scour-damage repairs along US 101 at the location on the outside of a meander bend in the channel-migration zone of the Hoh River near Forks, Washington. Four emergency projects that involved armoring the bank with large volumes of rock occurred at this location in the past few years, yet erosion continued and US 101 remained in imminent danger of being washed out.

Engineering analysis conducted by WSDOT indicated that relocation of the highway further from the channel-migration zone was economically infeasible. Therefore, bank-stabilization and river-deflection measures to protect the roadway were the only viable option. Because the "traditional" repairs were not effective, WSDOT developed an alternative solution for the site using engineered logjams (ELJs) in place of armoring bank-stabilization methods. The project has the added benefits of restoring salmon habitat and proving that sustainable engineering is not only possible, but can at times provide the most practical long-term solution.

ELJs emulate historic conditions and natural processes to rehabilitate aquatic and riparian habitat; provide erosion control, flood diffusion, and grade control; and increase sediment retention. Engineered logjams are an emerging technology based upon the premise of applying rigorous scientific and engineering principles to the design and construction of structures to protect infrastructure in a manner that emulates natural systems.

The Hoh River engineered-logjam project is the largest engineered-logjam project in the Pacific Northwest, and possibly the world. A series of 12 mid-channel and bank structures were installed. This action was intended to deflect and diffuse river flows to reduce the erosive forces acting upon the bank adjacent to the highway, as well as provide greater separation of the river from the highway shoulder.

The mid-channel logjam structures each include more than 100 logs (many with rootwads) with key log diameters of 36 to 48 inches. The core of each structure consists of steel H-piles, 65 logs, and 2,200 tons of rock. Each mid-channel structure is approximately 30 feet in height, 75 feet wide, and 70 feet long, with approximately 15 feet of the structure buried below the riverbed level. The structures include several large protruding logs that are used to hold smaller racked logs in place forming irregular faces. Exterior racked logs and naturally accumulating woody debris are key for complex habitat formation.

The design life of the engineered logjam structures is expected to be a minimum of 50 years. These structures will provide stable hard points that deflect river flow and provide a medium for the growth of native vegetation on logjam islands in the channel, while emulating natural logjams in many pristine river reaches in the Pacific Northwest.

The project was designed in the spring of 2004 and constructed from early July through mid-October 2004. Significant difficulties with the temporary river diversion, water-quality maintenance, and fish handling occurred during construction. Although winter flows have been lower than normal in this first year, indications are that the structures are performing as desired.

Biographical Sketch: Carl Ward received a B.S. in wildlife and fisheries biology from the University of Vermont (1987). He has been the WSDOT regional biologist/Olympic Region biology program manager since 1991. He manages a staff of biologists and an environmental-restoration work crew. Ward leads the team in the identification and evaluation of the effects of transportation design, construction, and operations on plants, wildlife, fisheries, threatened and endangered species, and associated habitats. The team also prepares and implements restoration, enhancement, and replacement mitigation plans for unavoidable impacts to natural resources. Prior to joining WSDOT, Carl worked in the private environmental-consulting field for four years in the assessment of wetlands, fish, and wildlife.

ENVIRONMENTAL RETROFIT FOR HIGHWAYS: MAKING WILDLIFE A PRIORITY

Paul Wagner (Phone: 360-705-7406, Email: wagnerp@wsdot.wa.gov), Biology Branch Manager, Washington State Department of Transportation, Environmental Services, P.O. Box 47331, Olympia, WA 98504, Fax: 360-705-6833

Abstract

The environmental aspects of transportation projects have typically focused on the avoidance and minimization of impacts and compensatory mitigation for unavoidable impacts. Recently, progressive transportation agencies have been expanding beyond the primary focus of project effects and evolving toward a more thorough integration of environmental stewardship in their actions. Agencies are beginning to integrate environmental factors into transportation planning and are also providing environmental enhancements as part of projects when opportunities arise.

Transportation agencies have traditionally prioritized their work to meet the typical infrastructure needs for addressing deficiencies and making improvements for safety, capacity, and system efficiency, as well as upgrading aging facilities. Significant environmental improvements can be reached using good stewardship practices in planned transportation projects. However, sometimes areas with ecological needs do not coincide with areas needing transportation-infrastructure improvement.

How can transportation programs move beyond the project and permit perspective and work to address ongoing ecological issues and thus provide larger environmental gains?

One approach is the Environmental Retrofit Program developed by the Washington State Department of Transportation (WSDOT). This program is designed to identify environmental deficiencies within the highway system and address them both as parts of planned transportation projects and also as stand-alone environmental-retrofit projects. These stand-alone retrofit projects may be conducted not only where the transportation needs are currently satisfied, but where significant ecological impacts exist. The focus areas for this program are based on the ecological priorities, including fish-passage correction, stream-habitat restoration, and water-quality improvements.

An example of the benefits of this program can be seen in fish passage retrofit activities. Culverts at road crossings that block fish movement are recognized as a significant conservation issue, particularly for anadromous salmonids, many of which are listed under the federal Endangered Species Act (ESA).

Since 1991, the Fish Passage Retrofit Program has been managed cooperatively between WSDOT and the Washington State Department of Fish and Wildlife (WDFW). Over 5,000 stream crossings have been inspected on the state highway system. As a result, over 800 culverts have been identified that block significant habitat upstream and are targeted for correction. Over \$26 million has been invested in inventory, design, and construction for stand-alone retrofit projects that restore fish passage at 59 high-priority sites.

As a result, access to over 400 linear miles of salmonid habitat, once blocked, has been improved. This presentation will discuss how the program operates, as well as specific examples of the projects that have been implemented.

The main components for operating this program include: Definition of the problem and parameters; Field Inventory and survey; Statewide prioritization, based on ecological gain; Scoping of project corrections; Design development; Permitting; Construction; Monitoring; Research; and Coordination and partnerships

The concepts of this program are now being expanded to address other types of aquatic-habitat issues though identification of what is termed chronic environmental deficiencies (CED) and stormwater treatment needs. Future applications of this program are being developed to address priorities for terrestrial habitat-connectivity improvement. This is a successful program with tangible benefits on the ground that demonstrates how transportation agencies can play a meaningful role in ecological-restoration efforts.

Biographical Sketch: Paul Wagner is a wildlife biologist with over 20 years experience in the field, including work with red-wolf reintroduction in North Carolina and studies of seabirds in Alaska's Pribilof Islands and ice-age mammals in Arctic Alaska. He is currently the biology branch manager for the Washington State Department of Transportation and manages programs responsible for policy and interagency coordination related to wetlands, fish, wildlife, and habitat issues statewide. He has a B.S. degree in natural history from Juniata College and graduate coursework in salmon ecology at Evergreen State College. He has served on committees of the National Academies of Sciences, involved in assessing the ecological effects of roads, and has been a steering committee member of ICOET since 1998.

RESTORATION OF AQUATIC HABITAT AND FISH PASSAGE DEGRADED BY WIDENING OF INDIAN HIGHWAY 58 IN GARHWAL HIMALAYA

Ramesh C. Sharma (Phone: 91-1370-267740, Email: drrameshcsharma@yahoo.com), Professor and Head, Department of Environmental Sciences, H.N.B. Garhwal University, Post Box 67, Srinagar-Garhwal 246174, Uttaranchal, India

Abstract: Sustainable approaches to the construction and widening of roads and highways are essential to offset negative influences on aquatic habitat and fish passage in the fragile ecosystem of the Himalayan Mountains in northern India. Evidence is growing that the expanding, poorly designed network of roads and trails in mountain areas, without giving due considerations to natural processes such as geological processes and climatic severity, such as heavy monsoon precipitation, is a major cause of habitat fragmentation and degradation of both terrestrial and aquatic habitats.

These effects have been quantified for aquatic primary producers (periphyton), aquatic benthic insects, and Snow Trout, a Himalayan teleost (*Schizothorax richardsonii*, Gray; *Schizothoracichthys progastus*, McClelland) that dwells in the upper Ganges River, following Indian National Highway 58 (NH-58) in the mountain region of Garhwal Himalaya, India (latitude 29 degree 61 minutes -30 degree 28 minutes N; longitude 77 degree 49 minutes -80 degree 6 minutes E). Indian Highway 58 is one of the most important highways, is 300-km long, and passes along the Alaknanda River (230 km), which is one of the parent streams of the Ganges (70 km) in the fragile mountain ecosystem of Garhwal Himalaya of northern India. Keeping in mind the heavy traffic on the highway, a RS 450 million (US \$100 million) widening project was launched in 2001.

The widening of Highway 58 through massive cutting of mountain slopes, the disposal of tons of the cut material downhill into the waterways in an uncontrolled manner, and the improper water management of the slopes has resulted in intensive accumulation of soil and woody debris into the aquatic ecosystem from accelerated erosion, gullies, and landslides, resulting in drastic changes in the physico-chemical and biological profile of the aquatic habitat. Detrimental effects on transparency, current velocity, conductivity, bottom-substrate composition, dissolved oxygen, periphyton production, and the production of benthic insect communities have been documented. Feeding, spawning, and the passage of the Snow Trout cold-water fish have been degraded or destroyed.

Subsequent to the widening of Highway 58, the annual gross primary production (Pg) of periphyton declined from 8771 g C m⁻³yr⁻¹ (96.48 k cal m⁻³yr⁻¹) to a value of 5952 g C m⁻³yr⁻¹ (65.47 k cal m⁻³yr⁻¹), a 32-percent decrease in aquatic habitat. The maximum biomass (standing crop) of aquatic insects declined from a mean monthly biomass of 4.926 g m⁻² (February) to 1.848 g m⁻², a 62-percent decrease, and a minimum monthly mean biomass of 0.408 g m⁻² (August) to 0.126 g m⁻², a 69-percent decrease. Subsequent to widening of the highway, the standing crop estimate of Snow Trout declined from a maximum mean monthly biomass of 2.955 g m⁻² (February) to 1.201 g m⁻², a 59-percent decrease, and a minimum monthly mean biomass (August) of 0.244 g m⁻² to 0.082 g m⁻², a 66-percent decrease. Annual productivity of Snow Trout declined from 1.309 g m⁻² to 0.448 g m⁻², a 66-percent decrease.

This decline is believed to have been caused by increased turbidity accompanied by a decline in depth and dissolved oxygen, accumulation of fine silt and suspended solids, a decrease in primary productivity, a decrease in general benthic-aquatic insects productivity, depletion of the food supply, and loss of cover.

We have recommended the following measures to restore habitat quality and connectivity of Snow Trout:

- Stream restoration and stream bank stabilization using these structures:
 - Toe walls
 - Retaining walls
 - Stone layers
 - Stone arches
 - Terraces
- Bioengineering methods by
 - Planting fast-growing plant species in combination with wire netting, gravel mining, and dredging in the impacted sites
 - Protecting riparian vegetation
 - Monitoring of water quality
 - Enhancement of fish food reserves
 - Sustainable approaches to road construction and widening
 - Proper drainage of water-saturated mountain slopes and spring runoff during monsoon season (July-August)
 - Sealing of side drains against underground water penetration alongside endangered sections of the highway
 - Construction of check dams for protection of steep gullies and side erosion of the river bed

We also recommend establishment of a strong partnership among experienced, expert

- Geologists
- Civil engineers
- Structural engineers
- Environmental biologists

Introduction

It is undisputed that the existence of roads is an imperative requirement for mobility, accessibility, and smooth development in Himalayan Mountains. Due to the young geology of the Himalayas and instability of their slopes, the region is prone to recurrent and often devastating landslides triggered by construction and widening of roads and highways.

Garhwal Himalaya is an important part of Himalaya located in the state of Uttaranchal of North India. The major Indian rivers (Ganges and Yamuna) and their tributaries (Alaknanda, Bhagirathi, Bhilangana, Mandakini, Pindar, and Nayar) have their origin in Garhwal Himalaya. Most of the roads in Garhwal Himalaya have been constructed in the valleys along the rivers. Therefore, any activity related to the construction and widening of roads has detrimental effects on the aquatic ecosystem and the organisms dwelling in it. Geology and the fragile nature of the region make this relationship of roads and the aquatic ecosystem more vulnerable.

Evidence is growing that the expanding, poorly designed network of roads and trails in mountain areas, without giving due consideration to natural processes such as geological processes and climatic severity, such as heavy monsoon precipitation, is a major cause of habitat fragmentation and degradation of both terrestrial and aquatic habitats. Massive cutting of the mountain slopes and disposal of the cut material downhill in an uncontrolled manner, uncontrolled blasting of rock in large quantities for road cutting, and improper water management in mountain terraces has resulted in intensive soil loss from accelerated erosion, gullying, and landslides.

Therefore, sustainable approaches to the construction and widening of roads and highways are essential to offset negative influences on the aquatic habitat and fish passage of the fragile ecosystem of the Himalayan Mountains of Northern India. Considerable work has been done on the impact of the transportation network on aquatic ecosystems and fish life in America and Europe. However, except by Sharma (2003), no sincere effort has been made so far on the restoration of aquatic habitat and fish passage degraded by roads and highways in India.

The present paper attempts to provide manifestation of the negative impact of Highway 58 on water quality and to quantify the impact on primary production (periphyton), secondary production (aquatic insects), and production of Snow Trout, an important Himalayan teleost. For Snow Trout, several remedial measures for restoring the habitat quality and connectivity degraded by the widening of NH-58 in Garhwal Himalaya have been suggested and tried on many stretches of the Alaknanda River.

The Snow Trout is an important coldwater fish distributed along the Himalayas in India, Pakistan, Bhutan, Nepal and Bangladesh. It contributes more than 65 percent of the total fish catch of Garhwal Himalaya. This is an important indigenous teleost dwelling in snow-fed hill streams of the Himalayas. Members of the subfamily Cyprininae pertaining to the family Cyprinidae are commonly known as Snow Trout. Snow Trout comprise mainly two genera: Schizothorax with a suctorial lower lip and Schizothoraichthys with a non-suctorial lower lip. The most important species of Snow Trout dwelling in Garhwal Himalaya streams are *Schizothorax richardsonii* (Gray) and *Schizothoraichthys progastus* (McClelland). The fish weigh up to 2.5 kg with a maximum total length of 45 cm. Snow Trout are surface feeders. They are local migratory fish and prefer temperatures between 5° to 20° Celsius (Singh and Sharma 1998, Sharma 2003).

Materials and Methods

Physiography of the study area

The study area is located in the Garhwal Himalaya, an important zone of the Himalayas and a part of the new state of Uttaranchal of North India (latitude: 29 degrees 26 minutes -31 degrees 28 minutes N; longitude: 77 degrees 49 minutes -80 degrees 6 minutes E). It encompasses six districts (Dehradun, Tehri, Pauri, Uttarakashi, Chamoli, and Rudraprayag) and covers an area of 30,029 km².

The area is very rich in biodiversity (animal, plants, and microbes). The entire region of Garhwal Himalaya is bestowed with tremendous freshwater resources in terms of major fluvial systems of Ganges and Yamuna and their tributaries. Due to the rich freshwater resources, Garhwal Himalaya is known as the 'tower of freshwater resources.' Two major parent streams, the Alaknanda and Bhagirathi, form the Ganges after confluences at Deoprayag.

A thick network of roads and highways has been launched in the region to cater to the needs of the heavy influx of tourists. Most of the roads and highways in Garhwal Himalaya have been constructed in the river valleys along the rivers.

Geology of the study area

The study area is characterized by a flat-topped ridge, steep slopes, and a wide valley. The area is covered by three types of rocks of the upper Proterozoic to lower Paleozoic ages (Valdiya 1984). The area is represented by huge, thinly foliated, highly folded, fractured, and joined phyllite rock traversed by quartz veins and few basic intrusions in the form of a sill and dykes. The phyllite is called Pauri phyllite (Kumar and Agrawal 1975). Vertically folded, highly fractured, pinkish ripple, and current-bent quartzite rocks, intercalated with a massive intrusion of meta volcanic rocks, are under the Garhwal groups of rocks. The tectonic features generally control the landform of an area; slopes of a drainage pattern are more sensitive to recent neotectonic activities.

The wide valley of the Alaknanda River is characterized by the set of terraces formed by the river shifting and reducing the water discharge. The river flowing in the area was assumed to have heavy water discharge with laminar flow that reduced to its present level. Therefore, the sediments and load deposited along the riverside in the form of terraces. Most of the lowest terraces are in contact of the river.

The whole stretch of the Alaknanda River covers a distance of about 250 km and flows across the different litho-tectonic units of Garhwal Himalaya. The river can be conveniently divided into three zones: Mana to Vishnu Prayag

(highest gradient-I grade), Vishnu Prayag to Karanprag (moderate gradient-II grade), and Karanprayag to Devprayag (low gradient-III grade). National Highway 58 runs along the Alaknanda River from Badrinath to Devprayag (230 km) and Deoprayag to Rishikesh (70 km) along the Ganges.

Natural preconditions for road construction and widening in Garhwal Himalaya

Road construction and widening are very much dependent on the natural preconditions (climate, geology, topography, and environment) in mountainous areas. Favorable preconditions generally result in modest construction/widening volume per km, whereas unfavourable preconditions can bring enormous work volume and be very expensive. The climate of Garhwal Himalaya is mainly dependent on the altitude and varies from subtropical to alpine and temperate. The annual rainfall differs from place to place, ranging from less than 250 mm to 3500 mm. Most of the precipitation (80 percent) occurs during the monsoon period (July-August), creating tremendous problems for the road builders.

Garhwal Himalaya is affected by a constant tectonic uplifting accompanied by a down cutting of the river systems. The results of these natural forces are slopes which become steeper and steeper and therefore unstable. It is evident that such conditions make road widening a difficult task. The hilly belt of Garhwal Himalaya generally consists of rugged topography with a tremendous difference in elevation ranging from 350 m above mean sea level (m.s.l.) to 3,500 m above m.s.l. The resulting steep slopes are divided into many gullies and small valleys, and the valley floors are extremely narrow.

Such extreme conditions demand very careful road construction and widening activities. Forest and vegetation cover is a must for a balanced ecosystem. Depletion of forest resources by cutting of trees for firewood (the source of energy) and the extension of farmland into steep and unstable areas has made the entire mountain area of Garhwal Himalaya vulnerable. Such deforested and abandoned land has accelerated water runoff in volume and speed and is prone to slides. These four natural preconditions have a negative influence on road construction and widening in Garhwal Himalaya.

Salient features of the Indian Highway 58

Highway 58 is one of the most important highways and is 300 km long, passing along the Alaknanda River (230 km) and the Ganges (70 km) in the fragile mountain ecosystem of Garhwal Himalaya of northern India. NH-58 caters to the needs of heavy traffic (0.5 million people per year) and is used by people visiting the world-famous Indian shrines of Badrinath, Kedarnath, and Kemkunth Sahib, in addition to world-heritage sites (Nanda Devi Biosphere Reserve and Valley of Flowers).

Keeping in mind the heavy traffic on NH-58, in 2001 a project costing over Rs 450 million (US \$100 million) for widening the highway was launched. The basic objective of widening NH-58 is to make it double lanes for the smoother running of traffic. This project is expected to be completed in March 2007. The details of the widening work on different stretches of NH-58 are presented in Table 1.

Table 1. Details of widening works of NH-58.

| Stretch | km | Activity | Expected Completion |
|-------------------------------|----|------------------------------------|---------------------|
| Byasi-Kodiyala | 08 | Cutting Work | March 2006 |
| Bagwan-Srinagar-Srikot | 20 | Hotmix | April 2005 |
| Pharaso-Kaliasaur-Rudraprayag | 21 | Cutting Work | April 2005 |
| Bugwan-Rudraprayag | 60 | Construction of culvert and Hotmix | March 2006 |
| Gauchar-Karnaprayag | 32 | Cutting Work | March 2006 |

Methodology

Physico-chemical analysis of the water quality of the aquatic habitat of the Alaknanda River was made following the methods outlined in Wetzel and Likens (1991) and APHA (1998). Primary productivity of periphyton was determined by incubating substrates in a 1.93-liter molded-polystyrene chamber (Rodgers et al. 1978) for a four-hour incubation (0800-1200 hrs). A submersible pump (powered by an attached battery pack) supplied water circulation in the chamber. A variable resistor allowed variable flow in the chamber. Black-plastic tape was used to cover the dark (opaque) chambers. For the oxygen produced over a given time period, calculations were made after running modified Winkler's test on each of the samples.

Calculations for gross primary productivity (Pg) were made as follows:

Gross Primary Productivity (Pg): Total oxygen produced = Oxygen at end, light chamber not covered (minus) oxygen at end, black-taped chamber

The values obtained in mg. l-1 oxygen were converted to milligrams of carbon per cubic meter (mg C m-3) multiplying the value by 375.36 (Strickland and Parsons 1960). The values in mg Cm-3 can be converted to grams of dry weight by multiplying the milligrams of carbon by two and dividing by 1000. The values of dry weight were converted to calories of energy multiplied by 5.5 (Benton and Warner Jr. 1972).

The productivity of aquatic insects (secondary productivity) was determined by the biomass method (Winberg 1971, Downing and Rigler 1984). For the study of density, biomass, and production of the Snow Trout, the three small seine nets (TSSN) methods of Penczak and O'Hara (1983) were employed. The value for instantaneous growth (G) was estimated, when growth is considered to exponential:

$$G = \frac{\log_e \bar{w}_2 - \log_e \bar{w}_1}{t}$$

Where \bar{w}_1 and \bar{w}_2 = mean weight of the fish at times t1 and t2, respectively.

To estimate monthly production, mean biomass (\bar{w}) was multiplied by the instantaneous growth rate (G): P= $\bar{w} \times G$ (Chapman 1978). The annual production (g.m-2.yr-1) was estimated for Snow Trout.

Results and Discussion

Morphometric transformation of fish habitat

A large-scale morphometric transformation of the habitat of Snow Trout in a large section of the Alaknanda River has taken place due to widening activities on NH-58. As a consequence of these construction activities accompanying the road widening, a large stretch of the fluvial system has been transformed into a trench and a dammed pool of sluggish currents of water from rapids, cascades, and part of the high water from riffles. The other section of the river has been converted into narrow, turbulent, and turbid riffles from white and clear water pools as a result of the large scale of disturbances caused by disposal of tons of cut material downhill into the waterways of the Alaknanda River (figure 1).



Figure 1. Disposal of cut material downhill into the Alaknanda River caused by the widening activities on NH-58.

The composition of bottom substrates has been drastically altered by the widening activity of NH-58. Improper management of the slopes has resulted in intensive accumulation of soil, woody debris into the aquatic ecosystem from accelerated erosion, gullying, and landslides.

Degradation of physico-chemical aquatic environment

Degradation in the mean physico-chemical parameters of the aquatic environment of the Alaknanda River caused by the widening of NH-58 over a three-year period (January 2002-December 2004) is presented in Table 2.

Table 2. Degradation in the mean physico-chemical parameters of aquatic environment of the Alaknanda River caused by widening of NH-58 during a three-year period (January 2002-December 2004)

| Parameter | Standard Site (S1) (± SD) | Impacted Site (S2) (± SD) |
|--------------------------|----------------------------|----------------------------|
| Air temperature (°C) | 21.42 ± 6.50 | 21.78 ± 6.53 |
| Water temperature (°C) | 13.35 ± 2.51 | 14.02 ± 2.21 |
| Hydromedian depth (m) | 2.34 ± 1.34 | 1.51 ± 1.20 |
| Conductivity (µScm-1) | 80.56 ± 24.67 | 82.09 ± 25.18 |
| Relative humidity (%) | 46.21 ± 5.58 | 41.65 ± 5.75 |
| Water velocity (m.sec-1) | 1.375 ± 0.685 | 1.15 ± 0.702 |
| Turbidity (NTU) | 85.21 ± 78.37 | 121.65 ± 84.73 |
| Transparency (m) | 1.581 ± 0.637 | 1.20 ± 0.416 |
| Photoperiod (LH day-1) | 11.76 ± 1.20 | 11.76 ± 1.20 |
| TDS (x 102 mg.l-1) | 4.90 ± 4.80 | 5.94 ± 4.94 |

Table 2 (continued)

| Parameter | Standard Site (S1) (\pm SD) | Impacted Site (S2) (\pm SD) |
|------------------------------|-----------------------------------|-----------------------------------|
| pH | 7.45 \pm 0.05 | 7.65 \pm 0.07 |
| Dissolved oxygen (mg l-1) | 13.8 \pm 3.12 | 8.54 \pm 0.68 |
| Free carbon dioxide (mg l-1) | 0.92 \pm 0.31 | 1.01 \pm 0.16 |
| Total alkalinity (mg l-1) | 39.54 \pm 6.54 | 37.51 \pm 5.10 |
| Phosphates (mg l-1) | 0.030 \pm 0.011 | 0.036 \pm 0.012 |
| Nitrates (mg l-1) | 0.023 \pm 0.012 | 0.30 \pm 0.013 |
| Silicates (mg l-1) | 0.039 \pm 0.34 | 0.043 \pm 0.038 |
| Sulphates (mg l-1) | 1.576 \pm 0.486 | 1.545 \pm 0.612 |
| Chlorides (mg l-1) | 3.104 \pm 1.112 | 3.281 \pm 0.765 |

Analysis of the data revealed that a slight change in the water temperature in year 2004 (14.02 ± 2.21 °C) was noticed in comparison to the temperature recorded before the project (13.35 ± 1.34 °C). The drastic change in hydromedian depth (HMD) was recorded at the impacted site (1.50 ± 1.20 m) in comparison to the depth at the reference site (2.34 ± 1.34 m). Conductivity was also influenced from the natural condition (80.56 ± 24.67 μ mho cm-1) by the widening activities at the impacted site (28.09 ± 25.18 μ mho cm-1).

The water velocity has been altered to a great extent at the impacted zone (1.15 ± 0.072 m sec-1) versus the water velocity at the unaltered site (1.375 ± 0.685 m sec-1). A considerable change in the suspended material in the water at the impacted section (121 ± 84.73 NTU) was recorded versus the reference site (85.21 ± 78.37 NTU). A reduction in dissolved oxygen was also recorded at the impacted site (8.54 ± 0.68 mg l-1) versus the reference site (13.8 ± 0.12 mg l-1). A minor change in other chemical parameters (free CO₂, phosphates, nitrates, sulphates, chlorides, and silicates) was also noticed at the impacted zone of the Alaknanda River in comparison with the study made before the initiation of the NH-58 project.

Trophic depression in the aquatic environment

The biotic profile of the aquatic environments of the Alaknanda River is characterized by the presence of periphyton and macrophytes at the primary trophic levels and zooplankton and aquatic benthic insects at secondary trophic levels. These biotic components act as food for hill stream fishes. The natural composition of these organisms was also drastically influenced by the widening activities of NH-58. The percentage of aquatic insects was reduced from 50.83 percent to 30.06 percent over a period of three years (figure 2) as a consequence of the degradation of aquatic environment caused by the NH-58 widening activities.




Figure 2. Impact of widening activities of NH-58 on the percentage composition of aquatic organisms of the Alaknanda River, Garhwal Himalaya, over a period of three years.

The road-widening activities of the Alaknanda River drastically influenced the primary production of the aquatic environment contributed by aquatic plants. The annual gross primary production of periphyton was reduced from 8771 g C m-3 yr-1 (96.48 k. cal. m-3 yr-1) to a value of 5952 g C m-3 yr-1 (65.47 k. cal. m-3 yr-1), a 32-percent decrease in aquatic primary production over a three-year period (figures 3 and 4) The peak in primary production was recorded during November-December (winter months), when the transparency of the water was recorded to be very high.




Figure 3. Impact of NH-58 widening on the annual carbon value of aquatic environment of the Alaknanda River.




Figure 4. Impact of NH-58 widening on gross primary production (Pg) over a period of three years (32 percent decrease).

The maximum biomass (standing crop) of aquatic insects declined from a mean monthly biomass of 4.926 g m⁻² (February) to 1.848 g m⁻², a 62 percent decrease and a minimum mean monthly biomass 0.408 g m⁻² (August) to 0.126 g m⁻², a 69-percent decrease (table 3).

Table 3. Impact of widening of NH-58 on the mean monthly biomass (g m⁻²) of aquatic insects

| Month | 2002 (g m ⁻²) | 2004 (g m ⁻²) |
|-----------|---------------------------|---------------------------|
| January | 3.342 | 1.062 |
| February | 4.926 | 1.848 |
| March | 4.788 | 1.812 |
| April | 3.624 | 1.362 |
| May | 2.052 | 0.708 |
| June | 0.672 | 0.258 |
| July | 0.420 | 0.138 |
| August | 0.408 | 0.126 |
| September | 1.092 | 0.408 |
| October | 1.632 | 0.606 |
| November | 2.022 | 0.738 |
| December | 2.400 | 0.846 |

Impact of widening of NH-58 on the life of Snow Trout

Inundation of Spawning and Feeding Grounds of Snow Trout

The inundation of spawning and feeding grounds of Snow Trout inhabiting the Alaknanda River was observed at the impacted site of the river. As a result of the road-widening activities of NH-58 and a phenomenal change in turbidity and silting pattern, the failure of spawning or ineffective spawning of Snow Trout was observed. The presence of gravel, pebbles, sand, and bank-side vegetation is a prerequisite for Snow Trout to build their spawning nests (redds).

Choking of breeding grounds and fish passage

Environmental degradation brought about by intensified road-widening activities in the Alaknanda River catchment has adversely affected the local migratory fish species (*Schizothorax richardsonii*, Gray; *Schizothoraichthys progastus*, McClelland). Due to land slides, slope failures, sliding of the retaining wall, and disposal of tons of cut material downhill into the waterways, substantial morphometric transformations have resulted in the fish habitat that obstruct the free movement of Snow Trout into the breeding grounds. To spawn, both species of Snow Trout need clean, stable, and well-oxygenated gravel habitats, shaded with riparian vegetation. After the eggs are laid in the gravel, well-oxygenated water must pass over the eggs (Sharma 1991).

Impact on Production of Snow Trout

As a consequence of the massive scale of road-widening activities in the entire valley of Alaknanda, the Snow Trout (an important food fish of Indian Himalaya) is facing a lot of survival problems in the degraded and stressed habitats. Various stages of the life cycle (migration, spawning, incubation, and rearing) of Snow Trout are drastically influenced (figure 5).




Figure 5. Impact of widening of NH-58 on the annual production (g m⁻²yr⁻¹) of Snow Trout of the Alaknanda River.

Mountain ecosystems play a key role in providing forest cover, feeding perennial river systems, conserving genetic diversity, and providing an immense resource base for livelihood to local inhabitants. However, the mountain ecosystems are among the most fragile ecosystems in terms of susceptibility to natural and anthropogenic shocks. There has been a significant adverse impact on the mountain fluvial ecosystem caused by construction and widening of roads and highways. In order to protect and restore the aquatic ecosystem for fish survival, the complex inter-relationships between fish and their habitat must be understood.

All cold-water fish species, including Snow Trout dwelling in Himalayan water, need relatively unaltered or pristine freshwater habitats during part or all of their life-cycle stages. Fish migration, spawning, incubation, and rearing are examples of life-cycle stages.

The successful completion of each of these stages is dependent on one or more of the following environmental conditions of the fresh water habitat:

- Water temperature
- Depth
- Velocity
- Turbidity
- Dissolved oxygen
- Bottom substrates
- Cover
- Food supply

Different aquatic communities are broadly associated with habitat features based on the following (Wetzel 1983, Cole 1994):

- Water temperature
- Salinity
- pH
- Flow velocity
- Plant nutrients
- Bottom substrates
- Water clarity
- Dependability of oxygen concentration
- Concentration of toxic material (Wetzel 1983, Cole 1994)

Early work on the influence of inorganic sediment on aquatic life has been reviewed by Cordone and Kelley (1961). The effects of construction of the M11 motorway in Essex, Great Britain, were studied by Extence (1978). The macro-invertebrate communities above and below the entry of motorway run-off became progressively dissimilar over the study period. Certain groups (such as stone flies, may flies, and cased caddis flies) were largely absent at the outset. These studies show that the high suspended solids carried by runoff during civil-engineering operations can have a marked effect on the ecology of the receiving stream. Their long-term effects could, however, prove to be small since, once the works are completed and winter spates have carried the bulk of the material away, recolonization can occur from upstream.

This view finds support in the studies of Barton (1977) who noticed that the reduced fish population (24 to 10 kg. ha⁻¹) immediately below the site of highway construction returned to the original level after the work had been completed. Duvel et al. (1976) reported that modification of streams had a direct deleterious effect on the trout population, and large trout were denied suitable natural hiding places (holes, undercut, and bank vegetation).

The relationship between fish life and suspended solids was the first to be considered by the European Inland Fisheries Advisory Commission in their Technical Paper Series (EIFAC 1964). This relationship has since been reviewed by Alabaster (1972) and Alabaster and Lloyd (1980). Trout population in stream sections affected by high suspended solids had lower densities than in unaffected stretches (Scullion and Edwards 1980).

The long linear ecosystems (rivers and streams) are particularly vulnerable to fragmentation. There is a growing concern about the role of road crossings in altering habitat and disrupting river and stream continuity. Little consideration has been given to the ecosystem processes such as natural hydrology, sediment transport, fish and wildlife passage, or the movement of woody debris (Jackson 2003).

According to Mann and Penczak (1986), productivity levels of fish are affected by both biotic and abiotic influences, with the latter being of prime importance. Biotic variables (cover, food, and predation) have more influence in stable environments. Zaleswaki and Naiman (1985) demonstrated that abiotic factors (fluvial geomorphology, geology, and climate) were of primary importance in many situations. Zaleswaki et al. (1985) stressed that growth rates in headwaters (low-order streams) are primarily restricted by abiotic factors, especially temperature and trophic status. Egglisshaw (1970) demonstrated a relationship between fish production and availability of water flow and feeding sites. According to Power (1973), the presence of cover in the form of boulders and large stones greatly enhances the holding capacity of the river for fish and hence influences the production level. A deleterious effect of turbidity on fish production was noticed by Starrett and Fritz (1965). According to them, turbidity probably affects the procurement of food by sight-feeding fish. It also affects production of plankton and other food resources of fish.

The production level of fish is also dependent on light access and the amount and quality of autochthonous and allochthonous organic matter (Naiman 1983, Minshall et al. 1983) and temperature and its range (Elliot 1976, Edward et al. 1976). Thomas (1998) studied the effects of highways on western cold-water fisheries of North America. Highway network activities have an adverse impact on cold-water fish through loss of fish habitat, changes in habitat quality, isolation of populations, reduction, and invertebrate food supplies. Sheehy (2001) reported that roads are the major sources of sediment deposited in streams. This is especially critical when roads are adjacent to streams with sensitive species, where any sediment deposited into streams could have adverse effects.

Management of aquatic habitat may be as simple as adding a bottom structure, such as artificial reefs or spawning gravel for protective cover on reproduction (Kohler and Hubert 1993). Many degraded habitats can be cost-effectively aerated to increase oxygen concentration, fertilized to increase productivity of aquatic plants, or dredged to remove sediments. Habitat management integrates the management of entire watersheds. Sustaining an optimum balance of surface water and groundwater contributes to aquatic habitats by controlling erosion of sediments and nutrients.

The negative effects of environmental change in fish have been cumulative and interactive. Predictive understanding and effective management requires more holistic ecosystem approaches. The concept of recovery of ecosystem 'integrity' as the most appropriate means for obtaining optimum sustained benefits has gained considerable credence.

Restoration of Aquatic Habitat and Fish Passage

Efficiently protecting and restoring aquatic habitat and fish passage degraded by the transportation network is one of the most-needed management actions for natural resource managers throughout the world (Forman and Sperling 2002). Aquatic-habitat enhancement should be undertaken integrating natural channel-design techniques, aquatic-vegetation restoration techniques, and more traditional hydraulic and channel-design engineering practices (Welsche 1985; Nyman 1998, 2003). Successful treatments or techniques that directly protect or restore aquatic habitat impacted by roads are wildlife and fish passage improvement, channel and floodplain structure placements, and reconnecting water bodies (Doyle 2003). The utilization of passive treatment systems to mitigate the effects of acid mine drainage and acidic leachate discharge is a recent innovation in the restoration of aquatic ecosystems (Brookens et al. 2003).

Development of mountain-specific and sustainable infrastructures in mountainous areas requires multi-disciplinary inputs (Deoja 1994). Protecting and restoring aquatic habitat and fish passage of Snow Trout in the Alaknanda River along NH-58 in Garhwal Himalaya has become a priority. Therefore, the following measures have been recommended to restore habitat quality and connectivity for the Snow Trout.

Stream restoration and stream bank stabilization

Stream restoration and stream bank stabilization of the Alaknanda River can be made by improving the stability of a slope or to regaining stability of a slope after failure. Three different measures can be applied: improving the slope by making it as dry as possible (drainage system), supporting the slope by structures, or stabilizing by bioengineering methods. These three methods should be combined to achieve the optimum success. Stream-bank stabilization can be made through the protection structures (toe walls, retaining walls) to retain soil masses, other structures like stone layers, systems of stone arches, and terraces for preventing slope-surface erosion caused by the widening of NH-58. All these methods (improving the slope, support of the slope by structures, and bioengineering) have also been recommended by Schaffner (1987).

Bioengineering erosion-protection measures will be very effective in stabilizing unstable slopes at several locations on NH-58. Bioengineering measures consist mainly of planting fast-growing nonpalatable (*Alnus* spp.) species suitable to the climatic conditions of the site. Most important is the plant's capacity for deep rooting, thus increasing the soil surface and water-absorption power (drainage effect). There should be proper drainage of water-saturated slopes and spring runoff during monsoon seasons (July- August).

Another surface erosion protection measure is the combination of planting and mini-terrace construction out of wood. Finally, mini toe walls made of wood were also constructed. It has to be emphasized that areas with new plant cover have to be fenced off or watched by a watchman to avoid foraging by free-grazing animals, causing an eventual failure of the protective measure. The best method to prevent erosion on the uphill of NH-58 would be not to touch the mostly unstable slopes. They should be left uninhabited with their original forest cover.

Slopes drainage system

The activity of widening of NH-58 is a massive interference with the environment. Therefore, it should be handled with the utmost care. Thus 'kid-glove' approaches to road construction and widening should be applied, which include automatically the principle of preventing and minimizing erosion. Following this concept, slope failures have to be immediately repaired to prevent further extension and avoid the possibility that they become uncontrollable. Where the water runoff is not tightly checked, the system has to be improved to prevent creeps and slides. Early failures of the toe walls due to heavy precipitation during monsoon season (July-August) are very common in Garhwal Himalaya (figure 6). Therefore, several big culverts and check dams have been constructed for proper drainage throughout the length of NH-58 along the Alaknanda River in Garhwal Himalaya (figure 7).



Figure 6. Early failure of toe wall due to heavy precipitation during monsoon (July-August) on NH-58.



Figure 7. Construction of big culverts and check dams along NH-58 for proper drainage.

Sealing of side drains

Sealing of side drains should be made immediately against water penetration into the underground alongside endangered sections. Site drains should be discharged only into natural brooks, rivulets, and rivers. Steep gullies carrying increased water volume due to road-water discharge should be protected by check dams as far down as necessary to avoid depth and slide erosion of the river bed.

Gravel mining and dredging in the impacted sites

Fine silt and suspended solids are accumulated in the riverine ecosystem of the Alaknanda River. Snow Trout have difficulty in respiring, and their eggs are smothered. Turbidity reduces plant productivity to the extent that photosynthesis is impaired by reduced sunlight. Recreational opportunities are also lost.

Sediments usually refer to soil particles that enter the water column from eroding land. Sediments consist of particles of all sizes, including fine clay particles, silt sand and gravel. Suspended sediments can be traced to the road-construction source. Restoration of the impaired ecosystem of the Alaknanda River caused by the activities of widening NH-58 can be done in different ways to improve the aquatic habitat. Dredging and gravel mining are among the important ways to restore the impaired ecosystem. Dredging involves widening and/or deepening river channels to facilitate migration of fish. Dredging also maintains the flow of water and prevents clogging caused by silt.

Protecting the riparian vegetation

Riparian vegetation, stream bank morphology, overhanging vegetation, undercut banks, aquatic vegetation, and deep-water pools of the Alaknanda River are drastically altered by the debris generated by the widening of NH-58. These natural structures provide adult and juvenile cold-water fish with shade, resting areas, and protection from predation.

Riparian management is extremely critical for fish and wildlife populations (Thomas 1986). Riparian vegetation provides habitat cover for fishes and other wildlife, moderates stream temperatures, serves as a food source, and helps in stabilizing embankments (Welsch 1992). Riparian zones of the Alaknanda River received an inappropriate amount of the impact of the road cutting and widening of NH-58.

Riparian zones often are the most productive sites in a region because floodplains frequently have rich soils with plentiful moisture. They have a greater diversity of plant and animal species than adjoining ecosystems. Healthy riparian systems purify water as it moves through the vegetation by retaining sediments and by retaining water in aquifers beneath the floodplain. Riparian zones often are a diverse mix of wetland and upland vegetation, all of which are linked closely with the floodplain groundwater.

Maintaining proper amounts of herbaceous vegetation is a critical part of increasing sediment deposition and enhancing channel restoration in a hill stream system (Clary et al. 1996). Conversion of shrubland or woodland to herbaceous vegetation can greatly increase water yields. This is because grasses and forbs generally transpire much less water than do woody plants.

Recovery of ecosystem integrity

The negative effects of environmental change in fish habitats have been cumulative and interactive. Predictive understanding and more effective management require a more holistic ecosystem approach. Recovery of ecosystem 'integrity' is the most appropriate means for obtaining optimum sustained benefit and has gained considerable credence.

Monitoring of water quality

Monitoring of water quality of the Alaknanda River ecosystem is a prerequisite for maintaining the optimum physico-chemical conditions for Snow Trout. Monitoring of water quality will provide base data for improving the water quality for successful completion of different stages of life cycle (migration, spawning, incubation, rearing) of Snow Trout.

Sustainable approaches to the construction and widening of roads and highways

Development of mountain-specific and sustainable roads and highways in Garhwal Himalaya requires multidisciplinary inputs. An integrated development strategy based on geological, engineering, socioeconomic, and environmental factors is required for the construction and widening of roads and highways in mountainous areas. Mountain-specific design and approaches for construction of roads and their widening require access to comprehensive knowledge of geology, geotectonic, engineering, and economic analysis. Traditional civil engineers must be trained in mountain-specific skills.

Acknowledgements: I am pleased to acknowledge the financial support by the organizers of IOCET 2005 and the important role of Katie McDermott, Director, Technology Transfer, Center for Transportation and the Environment, North Carolina State University, Raleigh, North Carolina, in enabling me to contribute to IOCET 2005.

Biographical Sketch: Professor Sharma has a distinguished academic career. He graduated with zoology honors and obtained a master's degree in zoology in freshwater fishery biology. He obtained his doctorate (Ph.D.) in the environmental biology of fish and his doctor of science (D.Sc.) in environmental biology. For more than thirty years, he has had a wide experience teaching and researching environmental monitoring, bioenergetics, limnology, resource management, aquatic biodiversity, hyporheic biodiversity, microbial diversity, and transportation and environmental issues in the Himalayas. More than 12 research projects have been completed on these aspects. Under his supervision, 18 doctoral-research students have earned doctoral degrees and seven more students are engaged in research. He has sufficient professional experience and exposure by way of visiting and working at different research laboratories in India and abroad

(Canada, the Czech Republic, Poland, Sweden, and the United States.). He has published more than 104 research articles in journals of international repute. He has received several awards and gold medals (NATCON Environment Gold Medal 2001, Zoological Society of India Gold Medal 2001, Environmentalist of the Year Award 2003, Recognition Award Gold Medal 2004, and Indira Gandhi National Environment Award 2005). He is a fellow of many national and international societies. Currently, he is the professor and chairman of the Department of Environmental Sciences, H. N. B. Garhwal University, Srinagar-Garhwal, Uttaranchal, India.

References

- Alabaster, J. S. and R. Lloyd. 1980. *Water Quality Criteria for Freshwater Fisheries*. London: Butterworths.
- Alabaster, J. S. 1972. Suspended solids and fisheries. *Proc. Royal Soc. London*, 180: 395-406.
- Anon., 2001. National Bureau of Fish Genetic Resources, Annual Report. Lucknow.
- APHA, AWWA-WPCF. 1998. *Standard Methods of the Examination of Water, Sewage and Industrial Waste*. American Public Health Association, Washington, D.C.
- Barton, B. A. 1977. Short-term effect of highway construction on the limnology of a small stream of southern Ontario. *Freshw. Biol.* 7: 99-108.
- Benton, A. and W. E. Warner Jr. 1972. *Manual of Field Biology and Ecology*. Burgess, Minnesota.
- Brookens, A, T. Schmidt, R. Morgan, M. Kline, K. Kline, and D. Gates. 2003. Restoration of an upper headwaters coldwater ecosystem in western Maryland utilizing passive treatment technologies. *ICOET 2003 Proceedings*.
- Chapman, D. W. 1978. *Production in Fish Population: Ecology of Freshwater Fish Production* (S. D. Gerking ed.). Blackwell, Oxford.
- Clary, W.P., C. I. Thornton, and S. R. Abt. 1996. Riparian stubble height and recovery of degraded streambanks. *Rangelands* 18: 137-140.
- Clinton, B. D. and J. M. Vose. 2002. Differences in surface water quality draining four roads surface types in the southern Appalachians. *Southern J. Applied Forestry*. 27(2): 100-106.
- Cole, G. A. 1994. *Text Book of Limnology*. Waveland, Prospect Heights, New York.
- Cordone, A. J. and D. W. Kelly. 1961. The influence of inorganic sediments on the aquatic life of stream. *Calif. Fish. Game* 47: 189-228.
- Deoja, B. B. 1994. Sustainable approaches to the construction of roads and other infrastructure in Hindu Kush Himalayas. *ICIMOD Occasional Paper No. 24*: 1-69.
- Downing, J. A. and F. H. Rigler. 1984. *A Manual on Methods for the Assessment of Secondary Productivity in Freshwater* (2nd ed.) IBP Handbook No. 17. Blackwell.
- Doyle, J. 2003. Protecting and restoring aquatic habitat connectivity along forest highways and low-volume roads. *ICOET 2003 Proceedings*.
- Duvel, W. A. et al. 1976. Environmental impact of stream channelization. *Wat. Resour. Bull.* 12: 799-812.
- Edwards, R.W., J.W. Densem, and P. H. Russel. 1976. An assessment of the importance of temperature as a factor controlling the growth rate of brown trout in streams. *J. Animal Ecol.* 45: 923-948.
- Egglishaw, H. J. 1970. Production of salmon and trout in a stream in Scotland. *J. Fish. Biol.* 2: 117-1436.
- Elliott, J. M. 1976. Energetics of feeding, metabolism and growth of brown trout, salmon trutta L. in relation to body weight, water temperature and ration size. *J. Anim. Ecol.* 45: 1-48.
- Extence, C. A. 1978. Effects of motorways construction on an urban stream. *Environ. Pollut.* 17: 245-262.
- Forman, R. T. T. and D. Sperling. 2002. *Road Ecology: Science and Solutions*. Island Press, Washington, D. C.
- Gaardner, J. and H. H. Gran. 1927. Investigation on the production of plankton in the Oslofjord. *Rapp. Proc. Vervaux Reunions Conseil Intern. Exploration. Mer.* 42: 1-48.
- Jackson S. 2003. Design and construction of aquatic organism passage at road-stream crossings: ecological consideration in the design of river and stream crossings. *ICOET 2003 Proceedings*.
- Kohlar, C. C. and W. A. Hubert (eds.). 1993. *Inland Fisheries Management in North America*. American Fishery Society, Bethesda, Maryland.
- Kumar, G. and N. C. Agarwal. 1975. Geology of the Srinagar-Nandprayag area, Alaknanda Valley. *Himalayan Geology*. 5: 29-59.
- Mann, R. H. K. and T. Penczak. 1986. Fish production in rivers: reviews. *Pol. Arch. Hydrobiol.* 33(3-4): 233-247.
- Minsall, G. W., R. C. Peterson, K. W. Cummins, J. L. Brit, J. R. Sedell, C. E. Cushing, and R. L. Vannote. 1983. Interbiome comparison of stream ecosystem dynamics. *Ecological Monograph* 53: 1-25.
- Morita, K. and S. Yamamoto. 2002. Effects of habitat fragmentation by damming on the persistence of stream-dwelling Charr population. *Conservation Biology* 16(5): 1318-1323.
- Naiman, R.J. 1983. Annual pattern and spatial distribution of aquatic oxygen metabolism in boreal forest watersheds. *Ecolo. Monogr.* 53: 73-94.
- Nyman, D. C. 2003. Aquatic habitat enhancement for Mad River and Beaver Pond Brook in conjunction with the reconstruction of I-82, Waterbury, Connecticut. *ICOET 2003 Proceedings*.
- Nyman, D. C. 1998. Restoring naturalized stream beds in an urban river channel. *Engineering Approaches to Ecosystem Restoration; Wetlands Engineering and River Restoration Conference* (Denver, Colorado, March 1998). American Society of Civil Engineers, Reston, Virginia.
- Penczak, T and K. O'Hara, 1983. Catch-effort efficiency using three small seine nets. *Fish Mgmt.* 14: 83-92.
- Penczak, T. 1981. Ecological fish production in two small lowland rivers in Poland. *Oceologia* 48: 107-111.
- Power, G. 1973. Estimates of age, growth, standing crop, and production of salmonids in some north Norwegian rivers and streams. *Rep. Inst. Freshwat. Res. Drottninghom* 53: 78-111.

- Rodgers Jr., J. H., K. L. Dickson, and J. Cairns Jr. 1979. A review and analysis of some methods of periphyton. Wetzel, R. L. (ed.). *Methods and Measurements of Periphyton: A Review*, ASTM Spec. Tech. Publ., Philadelphia, 142-167.
- Schaffner, U. 1987. Road construction in the Nepal Himalaya: the experience from the Lamosangu-Jari Road Project. ICIMOD Occasional Paper No. 8: 1-67.
- Scullion, J. and R. W. Edwards. 1980. The effects of pollutants from the coal industry on the fish fauna of a small river in the South Wales Coalfield. *Environ. Pollut.* 21: 141-153.
- Sharma, R. C. 1991. Ecological Impact of Tehri Dam Construction on Dynamics of Biological Production in the River Bhagirathi of Garhwal Himalaya. D. Sc. Thesis, H. N. B. Garhwal University, Srinagar-Garhwal, India, 1-118.
- Sharma, R. C. 2003. Fish Diversity and their Ecological Status in Protected Areas of Uttaranchal. S. R. Verma (ed.). *Protected Habitats and Biodiversity*, Nature Conservators Publications, 9: 617-638.
- Sharma, R. C. 2003. Protection of an endangered fish Tor tor and Tor putitora population impacted by transportation network in the area of the Tehri Dam Project, Garhwal Himalaya, India. ICOET 2003 Proceedings.
- Sheehy, D.M. 2001. Reconditioning and stabilization of unpaved roads for reducing road maintenance and impacts on fisheries habitat. ICOET 2001 Proceedings.
- Singh, D. and R.C. Sharma. 1998. Biodiversity, ecological status and conservation priority of the fish of the River Alaknanda, a parent stream of the River Ganges (India). *Aquatic Conservation: Marine and Freshwater Ecosystems* 8: 761-722.
- Starret, W. C. and A. W. Fritz. 1965. Biological investigation of the fishes of Lake Chautauqua, Illinois. *Illinois Natural History Survey Bulletin*. 29 (1): 1-104.
- Strickland, J. D. H. and T. R. Parsons. 1968. *A Practical Handbook of Seawater Analysis*. Bulletin of Fisheries Research Board, Canada.
- Thomas, A. E. 1998. Effects of highways on western cold-water fisheries. ICOWET 1998 Proceedings.
- United States General Accounting Office. 2001. Restoring fish passage through culverts on Forest Service and BLM lands in Oregon and Washington could take decades. GAO-02-136, Washington, D.C.
- Valdiya, K. S. 1984. *Environmental Geology: Indian Context*. Tata McGraw-Hill Pub., New Delhi.
- Winberg, G. G. (ed.). 1971. *Methods for the Estimation of Production of Aquatic Animals* (translated by A. Duncan), Academic Press, London.
- Welsch, D. J. 1992. *Riparian Forest Buffers: Function and Design for Protection and Enhancement of Water Resources*. U.S. Department of Agriculture, Forest Service, Radnor, Pennsylvania.
- Welsche, T. A. 1985. *Streams Channels Modifications and Reclamation Structures to Enhance Fish Habitat. The Restoration of Rivers and Streams: Theories and Experience*. J. A. Gore, ed. Butterworth, Boston.
- Wetzel, R. G. 1983. *Limnology*. Saunders, New York.
- Wetzel, R.G. and G. E. Likens. 1991. *Limnological Analyses*. Springer-Verlag, New York.
- Zalewski, M. and R. J. Naiman. 1985. *The Regulation of Riverine Fish Communities by a Continuum of Abiotic-Biotic Factors. Habitat Modification and Freshwater Fisheries*. (J. S. Alebaster ed.) Butterworth, London.
- Zalewski, M., P. Frankiewicz, and B. Brewinska. 1985. Factors limiting growth and survival of brown trout salmon trutta. *J. Fish. Biol.* 27(A): 59-73.

THE ROLE OF GEOMORPHIC RIVER REACH ASSESSMENTS IN DEVELOPING ENVIRONMENTALLY BENEFICIAL HIGHWAY-PROTECTION MEASURES

Jennifer Black Goldsmith (Phone: 206-441-9080, Email: jgoldsmith@herrerainc.com), Senior Scientist, and **Tim Abbe** (Phone: 206-441-9080, Email: tabbe@herrerainc.com), Director of River Science and Geomorphology, Herrera Environmental Consultants, Seattle, Washington 98121, Fax: 206-441-9108

Jim Park (Phone: 360-705-7415, Email: parkj@wsdot.wa.gov), Floodplain Management Specialist, Washington State Department of Transportation, P.O. Box 47331, Olympia, WA 98504-7331

Abstract

Historic highway placement within river valleys has commonly occurred within flood and erosion hazard areas. Traditional maintenance of highways and other infrastructure in these environments can be costly, result in significant environmental impacts, and exaggerate risk elsewhere. Many rivers are subject to frequent changes in position as they migrate within their valleys. This channel migration is not limited to low-lying land subject to frequent flooding, but can consume new areas where the river has not historically been.

Changes in channel geometry alter flow conditions that can lead to either degradation (down-cutting) or aggradation of the river. Degradation can undermine road grades and bridge abutments and piers. Aggradation can increase flood frequency. Chronic maintenance and emergency repair are expensive and often do not address the source of the problem, but rather address the effect the flooding and erosion is having on the highway and related infrastructure. Furthermore, these measures rarely address impacts to habitat or how habitat can be improved from a proposed project.

Conducting a “geomorphic reach assessment” of a river’s processes and dynamics can be a valuable management tool for highway maintenance and operations managers to better understand why maintenance measures are chronically failing and to minimize emergency response by assessing potential near-term river hazards that may pose a threat to a highway and infrastructure. Geomorphic assessments evaluate historic channel dynamics, current river conditions, and hydrologic characteristics of the river system.

These assessments can also include conceptual designs and recommendations describing how to protect the highway from flooding and erosion, as well as improve existing habitat that may have been historically compromised because of highway placement and maintenance.

The results of a geomorphic assessment provide useful scientific information that is used in developing effective design solutions that address the flooding and erosion problems associated with a highway in a manner that does not compromise habitat, but instead actually improves current habitat conditions.

One such emerging technology that was developed in the Pacific Northwest is the use of “engineered logjams” for highway and infrastructure protection with the secondary benefit of improving aquatic habitat. Logjams can increase pool frequency, channel length, and riparian cover, as well as provide necessary bank protection for highways located along actively eroding banks. However, these technologies reintroduce natural complexity and variability to the river system. An analysis of how these structures could potentially alter flooding and erosion within a reach needs to be assessed for individual site scenarios.

We present several examples of reach assessments conducted for the Washington Department of Transportation to provide a better understanding of highway segments with chronic problems and outline better long-term maintenance strategies that enhance habitat recovery. This approach was utilized in the implementation of a complex engineered logjam (ELJ) project that has successfully protected U.S. Highway 101 and created valuable new aquatic habitat in the Hoh River of western Washington.

Biographical Sketches: Jennifer Black Goldsmith is a senior scientist with Herrera Environmental Consultants in Seattle. Ms. Goldsmith has 14 years of experience conducting natural-resource assessments throughout the Pacific Northwest. Her professional expertise includes water resources, water quality, geomorphology, and forestry. Ms. Goldsmith has extensive experience preparing water-resource analysis documentation for a variety of environmental impact statements, reach analysis, environmental assessments, and permit applications for a variety of projects.

Timothy B. Abbe, Ph.D., L.E.G., L.H.G., director of River Science and Geomorphology, Herrera Environmental Consultants. Tim Abbe has 17 years of experience in geology, geomorphology, environmental restoration, applying engineering principles in environmental project design, and solving problems in urban fluvial and coastal environments. He has pioneered the development of engineered logjams, which are artificial structures that emulate naturally occurring stream structures to achieve particular purposes (e.g., bank protection, grade control, and sediment trapping). His work on engineered logjams has offered new technology to professionals who must comply with environmental regulations while solving traditional problems such as runoff and bank erosion.